

RESEARCH ARTICLE

Open Access



Effects of state-wide implementation of the Los Angeles Motor Scale for triage of stroke patients in clinical practice

Stefanie Behnke¹, Thomas Schlechtriemen², Andreas Binder³, Monika Bachhuber¹, Mark Becker², Benedikt Trauth¹, Martin Lesmeister¹, Elmar Spüntrup⁴, Silke Walter¹, Lukas Hoor², Andreas Ragoschke-Schumm¹, Fatma Merzou¹, Luca Tarantini¹, Thomas Bertsch⁵, Jürgen Guldner⁶, Achim Magull-Seltenreich⁶, Frank Maier⁷, Christoph Massing⁷, Volkmar Fischer⁸, Michael Gawlitza⁹, Katrin Donnevert⁹, Hans-Michael Lamberty¹⁰, Stefan Jung¹¹, Matthias Strittmatter¹², Silke Tonner¹², Johannes Schuler¹³, Robert Liszka¹³, Stefan Wagenpfeil¹⁴, Iris Q. Grunwald^{15,16}, Wolfgang Reith¹⁷ and Klaus Fassbender^{1*} 

Abstract

Background: The prehospital identification of stroke patients with large-vessel occlusion (LVO), that should be immediately transported to a thrombectomy capable centre is an unsolved problem. Our aim was to determine whether implementation of a state-wide standard operating procedure (SOP) using the Los Angeles Motor Scale (LAMS) is feasible and enables correct triage of stroke patients to hospitals offering (comprehensive stroke centres, CSCs) or not offering (primary stroke centres, PSCs) thrombectomy.

Methods: Prospective study involving all patients with suspected acute stroke treated in a 4-month period in a state-wide network of all stroke-treating hospitals (eight PSCs and two CSCs). Primary endpoint was accuracy of the triage SOP in correctly transferring patients to CSCs or PSCs. Additional endpoints included the number of secondary transfers, the accuracy of the LAMS for detection of LVO, apart from stroke management metrics.

Results: In 1123 patients, use of a triage SOP based on the LAMS allowed triage decisions according to LVO status with a sensitivity of 69.2% (95% confidence interval (95%-CI): 59.0–79.5%) and a specificity of 84.9% (95%-CI: 82.6–87.3%). This was more favourable than the conventional approach of transferring every patient to the nearest stroke-treating hospital, as determined by geocoding for each patient (sensitivity, 17.9% (95%-CI: 9.4–26.5%); specificity, 100% (95%-CI: 100–100%)). Secondary transfers were required for 14 of the 78 (17.9%) LVO patients. Regarding the score itself, LAMS detected LVO with a sensitivity of 67.5% (95%-CI: 57.1–78.0%) and a specificity of 83.5% (95%-CI: 81.0–86.0%).

Conclusions: State-wide implementation of a triage SOP requesting use of the LAMS tool is feasible and improves triage decision-making in acute stroke regarding the most appropriate target hospital.

Keywords: Prehospital, Large-vessel occlusion, Emergency medical services, Triage, Preclinical scale, Thrombectomy

* Correspondence: Klaus.Fassbender@uks.eu

¹Department of Neurology, Saarland University Medical Center, Kirrberger St. Bldg. 90, 66421 Homburg, Germany
Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Background

Stroke is a frequent cause of disability and death [1] with important medical and economic implications. When stroke is caused by large-vessel occlusion (LVO), trials provide compelling evidence that mechanical thrombectomy (MT) rather than medical treatment alone is most effective. Even so, although an estimated 10 to 25% of patients with stroke have LVO [2, 3], only a small minority (fewer than 7%) are treated with MT [2, 4, 5]. A main reason for this difference is that MT is not available at the many primary stroke centres (PSCs) but is offered only by a few specialized stroke centres (comprehensive stroke centres, CSCs).

Currently, stroke-management guidelines recommend the transport of all patients to the nearest stroke-ready hospital [6], which in most cases is a PSC not offering MT. Therefore, patients with LVO may secondarily be transferred to a CSC for thrombectomy. Importantly, compared with direct referral to a CSC, such interhospital transfers cause pronounced treatment delays ranging from 96 min to 111 min for patients with LVO [7–10]. Consistent with the “time is brain” concept [8], these delays significantly worsen clinical outcomes [9, 11]. On the other hand, not all patients with stroke should be transferred to CSCs because, apart from overwhelming already strained accident and emergency departments, bypassing PSCs could delay the administration of intravenous thrombolysis for most patients.

LVO is mostly associated with more severe stroke symptoms. Therefore, researchers have proposed the use in the field of stroke severity scales aimed at prehospitally detecting LVO; patients with LVO could then profit from direct transfer to a CSC. Recently, many LVO scales, such as the Los Angeles Motor Scale (LAMS) [12], the Rapid Arterial Occlusion Evaluation Scale (RACE) [13–15] or the A2L2 test (A, arm; L, leg) [16] have been proposed and studied by EMS in regard to their accuracy of detecting LVO.

Because the LAMS, apart from belonging to the most predictive of those scales [17, 18], requires assessment of only three motor symptoms (facial paresis, arm strength, and grip strength) [12, 19], it appears most suitable for implementation in prehospital emergency care protocols.

However, information on the effects of implementation of the LAMS for real-life triage of stroke patients is still missing. The aim of this study is to explore the effects of a state-wide triage standard operating procedure (SOP) requesting performance of the LAMS for triage decision-making in regard to the most appropriate target hospital.

Methods

Patients and study design

This prospective multicentre study, coordinated by the University of the Saarland, Germany, was opened on

March 1, 2018, and terminated on June 30, 2018. The trial was conducted in the federal state of Saarland in Germany, a mixed urban and rural state with an area of 1004 sq. mi (2571 km²) and approximately 992,000 inhabitants, in the context of a state-wide network of all certified stroke centres (eight PSCs, two CSCs). All emergency calls in this state are evaluated by a single dispatch centre. Inclusion criteria were suspicion of acute stroke by the EMS personnel on-scene, age of at least 18 years, and willingness to participate. Exclusion criteria were referral modes other than via EMS and critical illness requiring immediate transfer to the nearest intensive care unit.

Triage SOP based on the LAMS instrument

The stroke triage SOP was a rule for all EMS stations in the entire state. In this SOP, the following factors were considered in triage decision-making: (1) LAMS score (cut-off score of 4) [19]; (2) symptom onset times of 8 h or less or presence of “wake-up” stroke; and (3) quality-of-life aspects, such as severe comorbid conditions and severe prestroke dependency [20].

The triage protocol was set in operation by the central EMS coordinating authority of the state, the Zweckverband für Rettungsdienst und Feuerwehralarmierung, Saar, on May 30, 2015 (VAW MED-012) [20], and was accompanied by state-wide structured training sessions held approximately every 3 months for EMS personnel in the field and every 6 months for dispatch centre personnel in the context of the 30 educational sessions per year required for EMS personnel in the field. Moreover, the protocol was a component of the educational curriculum of the state’s EMS school.

Endpoints

The primary endpoint was accuracy of the triage SOP in triaging stroke patients to the appropriate target hospital, PSC versus CSC. Correct triage decision was defined as a decision to transport patients with LVO to the nearest CSC and to transport patients without LVO to the nearest PSC (or to a CSC, if this was the nearest stroke-treating hospital). LVO was defined as occlusion of the intracranial internal carotid artery, the proximal (M1) segments of the middle cerebral artery, or the basilar artery.

Secondary endpoints were performance of the triage protocol with regard to either LVO or intracranial haemorrhage (ICH), the number of secondary interhospital transfers, and the sensitivity and specificity of the LAMS in detection of LVO itself. Documented stroke management metrics included times from call to (1) “on scene”, (2) hospital admission (“door”), (3) first neurologist contact, (4) start of non-contrast imaging, (5) start of vascular imaging (in case of ischaemic stroke), (6)

needle, (7) door of CSC after secondary transfer, and (8) groin puncture. To determine short-term outcomes, we assessed mRS scores at discharge and mortality rates.

Geocoding

For analysis of the potentially alternative conventional pathway of the guideline-recommended transfer to the nearest stroke-treating hospital, geocoding was performed for each patient based on postal code, as previously described [21]. This approach allowed exact identification of the level of care (PSC or CSC) offered by the nearest stroke-treating hospital to which the individual patient with stroke would normally have been transferred.

Ethical aspects

The study protocol, the informed consent document, and the subject information document were approved by the Ethics Committee of the Medical Association of the Saarland, Germany (AZ-257/17). Informed consent was obtained from all patients or their legal representatives and was documented in the receiving hospitals. The otherwise separated datasets of the EMS and of the hospital were pseudonymized and linked only by the information from (1) the target hospital, (2) the admission date, (3) the patient’s sex, and (4) the patient’s year of birth.

Statistical analyses

Results are reported according to Strengthening The Reporting of OBservational Studies in Epidemiology (STROBE) guidelines [22]. Group comparisons were analysed with the Mann-Whitney U Test or the Fisher exact test. Statistical analyses were performed with IBM SPSS Statistics for Windows, version 25.0.0.0 (IBM Corporation, Armonk, NY, USA).

Results

Demographic and medical characteristics

After the 4-month study period, the study was terminated with a total of 1123 enrolled patients. Exclusion of patients is specified in Fig. 1. Baseline demographic and medical characteristics of the patients are presented in Table 1. Of the 1123 patients, 644 (57.3%) had experienced an acute ischaemic stroke, 84 patients (7.5%) had experienced a haemorrhagic stroke, and 395 patients (35.2%) had experienced stroke mimics. Of the 644 patients with ischaemic stroke, 129 (20.0%) underwent thrombolysis; and 489 underwent vascular imaging, which showed that 78 (12.1%) had LVO; 53 of these 78 patients (67.9%) were treated with MT.

Feasibility and performance of the LAMS-based triage SOP in triage of stroke patients

This study demonstrated the feasibility of state-wide implementation of a LAMS-based triage SOP with high adherence. Only 60 of 1123 patients (5.3%) had no




Fig. 1 Strengthening The Reporting of Observational Studies in Epidemiology (STROBE) diagram illustrating the trial design and exclusion of patients from the study. Patients with suspected stroke were enrolled in the hospital after evaluation for inclusion and exclusion criteria and after informed consent had been obtained. For pseudonymization, separated data sets from the prehospital and in-hospital phases of acute stroke management were linked exclusively on the basis of information about the date of stroke, target hospital, sex, and the patient’s year of birth

Table 1 Demographic and medical characteristics of the study population

	Total (n = 1123)	Triage decision		Transport destination ^a	
		CSC (n = 188)	PSC (n = 935)	CSC (n = 424)	PSC (n = 699)
Demographic and prehospital data					
Age, years; median (IQR)	78 (68–85)	74 (66–83)	79 (68–85)	76 (66–84)	80 (69–85)
Male sex, n (%)	565 (50.3)	97 (51.6)	468 (50.1)	218 (51.4)	347 (49.6)
Symptom onset to call, min; median (IQR)	55 (11–353)	19 (6–126)	68 (14–406)	36 (8–233)	74 (15–426)
LAMS, n/total n (%)					
LAMS < 4	855/1123 (80.4)	49/855 (5.7)	806/855 (94.3)	269/855 (31.5)	586/855 (68.5)
LAMS ≥ 4	208/1123 (19.6)	135/208 (64.9)	73/208 (35.1)	140/208 (67.3)	68/208 (32.7)
NIHSS admission, median (IQR)					
All	4 (1–8)	10 (5–17)	3 (1–6)	5 (2–11)	3 (1–6)
Patients with strokes	4 (2–8)	8 (5–16)	3 (1–6)	5 (2–10)	3 (1–6)
Discharge diagnoses, n (%)					
Ischaemic stroke	644 (57.3)	119 (63.3)	525 (56.1)	254 (59.9)	390 (55.8)
LVO stroke	78 (6.9)	54 (28.7)	24 (2.6)	58 (13.7)	20 (2.9)
Intracranial haemorrhage	84 (7.5)	28 (14.9)	45 (4.8)	56 (13.2)	39 (5.6)
Stroke mimics	395 (35.2)	41 (21.8)	354 (37.9)	125 (29.5)	270 (38.6)
Epileptic seizure	85 (7.6)	21 (11.2)	64 (6.8)	38 (9.0)	47 (6.7)
Migraine	11 (1.0)	1 (0.5)	11 (1.2)	3 (0.7)	8 (1.1)
Vestibulopathy	31 (2.8)	0	31 (3.3)	9 (2.1)	22 (3.1)
Infection	52 (4.6)	4 (2.1)	48 (5.1)	19 (4.5)	33 (4.7)
Exsiccosis	17 (1.5)	0	17 (1.8)	4 (0.9)	13 (1.9)
Delirium	31 (2.8)	0	31 (3.3)	8 (1.9)	23 (3.3)
Hypertensive crisis	6 (0.5)	1 (0.5)	5 (0.5)	2 (0.5)	4 (0.6)
Peripheral facial palsy	11 (1.0)	0	11 (1.2)	6 (1.4)	5 (0.7)
Intoxication	6 (0.5)	1 (0.5)	5 (0.5)	2 (0.5)	4 (0.6)
Syncope	18 (1.6)	1 (0.5)	17 (1.8)	4 (0.9)	14 (2.0)
Other	86 (7.7)	12 (6.4)	74 (7.9)	31 (7.3)	55 (7.9)

CSC Comprehensive stroke centre, PSC Primary stroke centre, IQR Interquartile range, LAMS Los Angeles Motor Scale, NIHSS National Institutes of Health Stroke Scale, mRS Modified Rankin Scale, LVO Large-vessel occlusion

^a Including CSCs serving as PSCs if they are the closest stroke centre at all

documented LAMS score, and 941 of the 1123 patients (83.8%) were triaged according to the LAMS score (Table 1).

The triage SOP allowed triage to the appropriate hospital with a sensitivity of 69.2% (95% confidence interval (95%-CI): 59.0–79.5%) and a specificity of 84.9% (95%-CI: 82.6–87.3%) (Table 2). In contrast, if the current guidelines had been adhered to (transfer to the nearest stroke-treating hospital), patients with LVO would have been correctly triaged with a sensitivity of 17.9% (95%-CI: 9.4–26.5%) and a specificity of 100% (95%-CI: 100–100%).

Performance of the LAMS in identifying LVO

The LAMS itself, at a cut-off value of 4, exhibited a sensitivity of 67.5% (95%-CI: 57.1–78.0%) and a specificity of 83.5% (95%-CI: 81.0–86.0%) in detecting LVO

(Table 3). Moreover, this scale exhibited a sensitivity of 56.6% (95%-CI: 48.9–64.3%) and a specificity of 86.7% (95%-CI: 84.3–89.1%) in detecting either LVO or ICH (Table 3). The receiver operating characteristic (ROC) curves obtained with a wider range of LAMS cut-off scores are displayed in Fig. 2 and suggest that the cut-off value of 4 is appropriate.

Similar results were obtained when the evaluation included ICH as a condition for triage to a CSC (Table 3). In accordance with this high triage accuracy, only 14 of the 78 patients with LVO (17.9%) required secondary transfer.

Stroke management metrics and short-term outcomes

Stroke management metrics are displayed in Table 4. The median call-to-needle times for patients transferred to a PSC were shorter than those for patients transferred

Table 2 Performance of the triage SOP and modelled conventional care assessed via geocoding in patients with LVO and with LVO or ICH^a

	Triage SOP (n = 968)	Modelled conventional care ^b (n = 968)
LVO, n/total n (%)		
Sensitivity	54/78 (69.2; 59.0–79.5)	14/78 (17.9; 9.4–26.5)
Specificity	756/890 (84.9; 82.6–87.3)	890/890 (100; 100–100)
Positive Predictive Value	54/188 (28.7; 22.3–35.2)	14/14 (100; 100–100)
Negative Predictive Value	756/780 (96.9; 95.7–98.1)	890/954 (93.3; 91.7–94.9)
LVO or ICH, n/total n (%)		
Sensitivity	82/162 (50.6; 42.9–58.3)	38/162 (23.5; 16.9–30.0)
Specificity	700/806 (86.8; 84.5–89.2)	806/806 (100; 100–100)
Positive Predictive Value	82/188 (43.6; 36.5–50.7)	38/38 (100; 100–100)
Negative Predictive Value	700/780 (89.7; 87.6–91.9)	806/930 (86.7; 84.5–88.9)

SOP Standard operating procedure, LVO Large-vessel occlusion, ICH Intracranial haemorrhage

^a 60 of 1123 (5.3%) patients had no documented LAMS score, and 155 of 644 (24.1%) ischaemic stroke patients did not undergo vascular imaging; ^b Conventional care was the guideline-recommended transfer to the nearest stroke centre, as calculated via geocoding in each patient. CSCs served as PSCs if they were the nearest stroke centre

95% confidence intervals are reported in brackets

to a CSC, and, conversely, call-to-groin puncture times were longer for patients transferred to a PSC than for those transferred to a CSC (Table 4).

In agreement with their lower LAMS scores, patients admitted to PSCs had better discharge modified Rankin scale (mRS) scores (1; interquartile range (IQR), 0–3) than did those admitted to a CSC (mRS score, 2; IQR, 1–5; $p < 0.001$). In addition, mortality rates were lower for patients transferred to a PSC (22, 4.8%) than for those transferred to a CSC (41, 11.9%; $p < 0.05$). Similarly, among patients with a stroke diagnosis, those admitted to PSCs had better discharge mRS scores (1; IQR, 0–3) than did those admitted to a CSC (mRS score, 2; IQR, 1–5; $p < 0.001$); they also exhibited lower mortality rates (12 patients, 3.6%) than did those admitted to a CSC (22 patients, 8.4%; $p < 0.05$).

Discussion

The current guideline-recommended practice of transferring each patient to the nearest stroke-treating hospital, usually a hospital not offering thrombectomy, may delay

or even preclude thrombectomy for patients with LVO. The results show that state-wide implementation of an SOP requesting the use of LAMS for triage decisions is feasible with high adherence and that this intervention can improve triage decision-making in regard to the appropriate target hospital in clinical practice.

Our evaluation of the effects of the state-wide implementation of an EMS SOP based on the LAMS achieved a sensitivity of 69.2% and a specificity of 84.9% in transferring patients with LVO to the appropriate target hospital. In accordance with such a high accuracy of triage decisions, we observed a low rate of secondary transfers (only 17.9% of LVO patients). In contrast, the conventional, guideline-recommended approach of transferring every patient to the nearest stroke-treating hospital, as determined by geocoding for each of the patients, would have achieved a sensitivity of only 17.9% and a specificity of 100%, a finding supporting the benefit of using the LAMS-based SOP. (The high specificity achieved by the modelled conventional pathway can be explained by the fact that “non-LVO” patients would always arrive at the “correct” hospital: the

Table 3 Performance of the LAMS tool (cut-off value, ≥ 4) in identification of LVO and LVO or ICH

Variable, n/total n (%)	All patients ^a (n = 920)		Patients with stroke ^b (n = 547)	
	LVO	LVO or ICH	LVO	LVO or ICH
Sensitivity	52/77 (67.5; 57.1–78.0)	90/159 (56.6; 48.9–64.3)	52/77 (67.5; 57.1–78.0)	90/158 (57.0; 49.2–64.7)
Specificity	704/843 (83.5; 81.0–86.0)	660/761 (86.7; 84.3–89.1)	366/470 (77.9; 74.1–81.6)	323/389 (83.0; 79.3–86.8)
Positive predictive value	52/191 (27.2; 20.9–33.5)	90/191 (47.1; 40.0–54.2)	52/156 (33.3; 25.9–40.7)	90/156 (57.7; 49.9–65.4)
Negative predictive value	704/729 (96.6; 95.3–97.9)	660/729 (90.5; 88.4–92.7)	366/391 (93.6; 91.2–96.0)	323/391 (82.6; 78.9–86.4)

LAMS Los Angeles Motor Scale, LVO Large-vessel occlusion, ICH Intracranial haemorrhage; Independently of having received vascular imaging, patients with ICH or stroke mimics are classified as LVO-negative

^a 60 of 1123 (5.3%) patients had no documented LAMS score, and 155 of 644 (24.1%) ischaemic stroke patients did not undergo vascular imaging; ^b 38 of 728 (5.2%) patients had no documented LAMS score, and 155 of 644 (24.1%) ischaemic stroke patients did not undergo vascular imaging

95% confidence intervals are reported in brackets




Fig. 2 Receiver operating characteristic curves of the Los Angeles Motor Scale at various cut-off scores for diagnosing large-vessel occlusion (LVO) selectively (dashed line) with an area under the curve of 0.798, or for diagnosing LVO or intracerebral haemorrhage (solid line) with an area under the curve of 0.751. Diagonal segments are produced by ties. The dotted line depicts randomness

nearest one, regardless of the MT options available there.) Thus, this finding strongly suggests that implementation of a protocol using the LAMS for triage decision-making is superior to the conventional approach of transporting every patient to the nearest hospital.

Similar values were obtained when ICH was included as a “target condition” for triage to a CSC. Although increasing evidence indicates that the “time-is-brain” concept is also valid for patients with haemorrhagic stroke and that these patients could also benefit from rapid

Table 4 Stroke management metrics

	Total (n = 1077)	CSC (n = 400)	PSC (n = 677)	p-value
Stroke-management metrics, min; median (IQR)				
Time from call to				
On-scene	10 (8–13)	9 (7–12)	10 (8–13)	0.105
Door ^a	49 (41–59)	51 (41–61)	48 (40–59)	0.008
First contact to neurologist	51 (42–62)	51 (42–62)	52 (43–62)	0.739
Non-contrast imaging	75 (60–99)	73 (58–104)	76 (62–96)	0.698
Initial vascular imaging	75 (62–99)	74 (60–99)	76 (64–98)	0.005
Needle	87 (67–100)	93 (73–103)	79 (66–95)	0.017
Groin puncture	112 (95–135)	105 (88–121)	292 (230–515)	< 0.001
Time from door to				
Non-contrast imaging	23 (13–42)	18 (11–43)	26 (16–40)	< 0.001
Initial vascular imaging	21 (15–38)	19 (14–33)	27 (20–42)	< 0.001
Needle	35 (27–49)	40 (29–50)	32 (26–48)	0.054
Groin puncture	57 (48–85)	53 (47–74)	233 (172–446)	< 0.001

CSC Comprehensive stroke centre, PSC Primary stroke centre, IQR Interquartile range

^a Time to admission at the first receiving hospital

specialist treatment at a CSC [12, 23], evidence from randomized studies regarding primary transfer to CSCs is still scarce [24].

Regarding the LAMS itself, this instrument detected LVO with a sensitivity of 67.5% and a specificity of 83.5% and detected either LVO or ICH with a sensitivity of 56.6% and a specificity of 86.7%. Calculating the accuracy of a variety of alternative LAMS cut-off scores confirmed that the cut-off score of 4 is appropriate. Thus, these are within the range of results of most earlier validation studies on the LAMS. E.g., a subgroup of 94 patients from the FAST-MAG trial who underwent vascular imaging found that the LAMS detected LVO with a sensitivity of 76% and a specificity of 65% [12], and two European studies reporting sensitivities of 63 and 38%, and specificities of 84 and 93%, respectively [17, 18].

However, these earlier studies did not use their LAMS results for triaging patients in clinical practice. At the same time, the results also corroborate the conclusions of previous meta-analyses indicating that all of the LVO scales tested still miss a substantial proportion of LVOs [25, 26].

While the strengths of this study are the pseudonymized linkage of separated prehospital and in-hospital data sets, and the high degree of data completeness, one limitation of the study is the lack of a control group. The alternative of retrieving historical data before implementing the SOP may, however, have caused a bias by secular effects, and randomization was not possible in this state-wide EMS rule. However, geocoding allowed us to exactly identify the level of care offered by the nearest stroke-treating hospital in each patient if the conventional practice had been applied. Furthermore, in this real-life-study, vascular imaging has not been performed in all patients admitted to the ten hospitals, thus resulting in omission of 24% of stroke patients from evaluation. In the future, triage accuracy may be further enhanced by improved scales, by the inclusion of additional factors such as screening tools for treatment eligibility or for mimicking conditions [27], or by additional telemedical consultation with the CSC team [16].

Conclusion

The prehospital identification of stroke patients with LVO that should be directly transported to a thrombectomy capable centre is an unsolved medical problem. Here, we show for the, to our knowledge, first time that a state-wide EMS SOP for stroke management requesting determination of the LAMS score is feasible with high adherence and can indeed be beneficial in triage decision-making regarding the most appropriate target hospital.

Abbreviations

95%-CI: 95% confidence interval; CSC: Comprehensive stroke centre; EMS: Emergency medical service; ICH: Intracranial haemorrhage; IQR: Interquartile range; LAMS: Los Angeles Motor Scale; LVO: Large-vessel occlusion; mRS: Modified Rankin Scale for Neurologic Disability; MT: Mechanical thrombectomy; NIHSS: National Institutes of Health Stroke Scale; PSC: Primary stroke centre; ROC: Receiver operating characteristic; SOP: Standard operating procedure

Acknowledgements

We thank the Ministry for Health Care and Social Affairs, and we thank the involved EMS health care professionals and hospital physicians for their excellent cooperation.

Authors' contributions

Conceptualization, S.B., T.S., A.B. and K.F.; data curation, S.B.; formal analysis, B.T., M.L., L.T., T.B. and St.Wa.; investigation, E.S., Si.Wa., L.H., A.R.-S., Fa.Me., L.T., T.B., J.G., A.M.-S., Fr.Ma., C.M., V.F., M.G., K.D., H.-M.L., S.J., M.S., S.T., J.S., R.L., I.Q.G.; project administration, T.S.; supervision, S.B., T.S. and K.F.; validation, S.B., Mo.Ba., Ma.Be., B.T., M.L. and St.Wa.; writing—original draft preparation, K.F.; writing—review and editing, S.B. All authors read and approved the final manuscript.

Funding

This work was funded by the Ministry of Health Care and Social Affairs of the Saarland, Germany.

Availability of data and materials

The datasets used and analysed during the current study are available from the corresponding author upon reasonable request.

Declarations

Ethics approval and consent to participate

The study protocol, the informed consent document, and the subject information document were approved by the Ethics Committee of the Medical Association of the Saarland, Germany (AZ-257/17). Informed consent was obtained from all patients or their legal representatives and was documented in the receiving hospitals.

Consent for publication

Not applicable.

Competing interests

Prof. Klaus Fassbender received grants from the Ministry of Health Care and Social Affairs of the Saarland, Germany. Dr. Binder received personal fees from Pfizer, Bristol-Myers Squibb, Bayer, and Daiichi Sankyo.

Author details

¹Department of Neurology, Saarland University Medical Center, Kirrberger St. Bldg. 90, 66421 Homburg, Germany. ²Zweckverband für Rettungsdienst und Feuerwehralarmierung Saar, Bexbach, Germany. ³Department of Neurology, Klinikum Saarbrücken, Saarbrücken, Germany. ⁴Department of Radiology, Klinikum Saarbrücken, Saarbrücken, Germany. ⁵Institute of Clinical Chemistry, Laboratory Medicine and Transfusion Medicine, Paracelsus Medical University, Nuremberg, Germany. ⁶Department of Neurology, Knappschaftsklinikum Saar, Püttlingen, Germany. ⁷Department of Neurology, Caritas-Klinikum Saarbrücken St. Theresia, Saarbrücken, Germany. ⁸Department of Neurology, Diakonie Klinikum Neunkirchen, Neunkirchen/Saar, Germany. ⁹Department of Neurology, Knappschaftsklinikum Saar, Sulzbach, Germany. ¹⁰Department of Neurology, DRK Krankenhaus Saarlouis, Saarlouis, Germany. ¹¹Department of Neurology, Marienhaus Klinikum Saarlouis-Dillingen, Dillingen, Germany. ¹²Department of Neurology, SHG Klinikum Merzig, Merzig, Germany. ¹³Department of Neurology, Marienhaus Klinik St. Wendel, St. Wendel, Germany. ¹⁴Institute for Medical Biometry, Epidemiology and Medical Informatics, Saarland University, Campus Homburg, Homburg, Germany. ¹⁵Department of Neuroscience, Medical School, Anglia Ruskin University, Chelmsford, UK. ¹⁶Division of Imaging Science and Technology, School of Medicine, University of Dundee, Dundee, UK. ¹⁷Department of Neuroradiology, Saarland University Medical Center, Homburg, Germany.

Received: 23 April 2021 Accepted: 4 May 2021

Published online: 01 June 2021

References

- Feigin, V. L., Norrving, B., & Mensah, G. A. (2017). Global burden of stroke. *Circulation Research*, *120*(3), 439–448. <https://doi.org/10.1161/CIRCRESAHA.120.030841>.
- Rai, A. T., Seldon, A. E., Boo, S., Link, P. S., Domico, J. R., Tarabishy, A. R., ... Carpenter, J. S. (2017). A population-based incidence of acute large vessel occlusions and thrombectomy eligible patients indicates significant potential for growth of endovascular stroke therapy in the USA. *Journal of Neurointerventional Surgery*, *9*(8), 722–726. <https://doi.org/10.1136/neurintsurg-2016-012515>.
- Lakomkin, N., Dharmoon, M., Carroll, K., Singh, I. P., Tuhim, S., Lee, J., ... Mocco, J. (2019). Prevalence of large vessel occlusion in patients presenting with acute ischemic stroke: A 10-year systematic review of the literature. *Journal of Neurointerventional Surgery*, *11*(3), 241–245. <https://doi.org/10.1136/neurintsurg-2018-014239>.
- Adeoye, O., Albright, K. C., Carr, B. G., Wolff, C., Mullen, M. T., Abruzzo, T., ... Kleindorfer, D. (2014). Geographic access to acute stroke care in the United States. *Stroke*, *45*(10), 3019–3024. <https://doi.org/10.1161/STROKEAHA.114.006293>.
- Weber, R., Eyding, J., Kitzrow, M., Bartig, D., Weimar, C., Hacke, W., & Krogias, C. (2019). Distribution and evolution of acute interventional ischemic stroke treatment in Germany from 2010 to 2016. *Neurological Research and Practice*, *1*(1), 4. <https://doi.org/10.1186/s42466-019-0010-8>.
- Powers, W. J., Rabinstein, A. A., Ackerson, T., Adeoye, O. M., Bambakidis, N. C., Becker, K., ... American Heart Association Stroke Council (2018). 2018 guidelines for the early management of patients with acute ischemic stroke: A guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*, *49*(3), e46–e110. <https://doi.org/10.1161/STR.0000000000000158>.
- Goyal, M., Jadhav, A. P., Bonafe, A., Diener, H., Mendes Pereira, V., Levy, E., ... For the SWIFT PRIME investigators (2016). Analysis of workflow and time to treatment and the effects on outcome in endovascular treatment of acute ischemic stroke: Results from the SWIFT PRIME randomized controlled trial. *Radiology*, *279*(3), 888–897. <https://doi.org/10.1148/radiol.2016160204>.
- Saver, J. L., Goyal, M., van der Lugt, A., Menon, B. K., Majoie, C. B., Dippel, D. W., ... Collaborators, H. (2016). Time to treatment with endovascular thrombectomy and outcomes from ischemic stroke: A meta-analysis. *JAMA*, *316*(12), 1279–1288. <https://doi.org/10.1001/jama.2016.13647>.
- Froehler, M. T., Saver, J. L., Zaidat, O. O., Jahan, R., Aziz-Sultan, M. A., Klucznik, R. P., ... Investigators, S. (2017). Interhospital transfer before thrombectomy is associated with delayed treatment and worse outcome in the STRATIS registry (systematic evaluation of patients treated with neurothrombectomy devices for acute ischemic stroke). *Circulation*, *136*(24), 2311–2321. <https://doi.org/10.1161/CIRCULATIONAHA.117.028920>.
- Prabhakaran, S., Ward, E., John, S., Lopes, D. K., Chen, M., Temes, R. E., ... Bleck, T. P. (2011). Transfer delay is a major factor limiting the use of intra-arterial treatment in acute ischemic stroke. *Stroke*, *42*(6), 1626–1630. <https://doi.org/10.1161/STROKEAHA.110.609750>.
- Saver, J. L., Goyal, M., Bonafe, A., Diener, H. C., Levy, E. I., Pereira, V. M., ... Swift Prime Investigators (2015). Stent-retriever thrombectomy after intravenous t-PA vs. t-PA alone in stroke. *The New England Journal of Medicine*, *372*(24), 2285–2295. <https://doi.org/10.1056/NEJMoa1415061>.
- Noorian, A. R., Sanossian, N., Shkirkova, K., Liebeskind, D. S., Eckstein, M., Stratton, S. J., ... Fast-Mag Trial Investigators and Coordinators (2018). Los Angeles Motor Scale to identify large vessel occlusion: Prehospital validation and comparison with other screens. *Stroke*, *49*(3), 565–572. <https://doi.org/10.1161/STROKEAHA.117.019228>.
- Pérez de la Ossa, N., Carrera, D., Gorchs, M., Querol, M., Millán, M., Gomis, M., ... Dávalos, A. (2014). Design and validation of a prehospital stroke scale to predict large arterial occlusion: The rapid arterial occlusion evaluation scale. *Stroke*, *45*(1), 87–91. <https://doi.org/10.1161/STROKEAHA.113.003071>.
- Carrera, D., Campbell, B. C., Cortés, J., Gorchs, M., Querol, M., Jiménez, X., ... Pérez de la Ossa, N. (2017). Predictive value of modifications of the prehospital rapid arterial occlusion evaluation scale for large vessel occlusion in patients with acute stroke. *Journal of Stroke and Cerebrovascular Diseases*, *26*(1), 74–77. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2016.08.032>.
- Zaidi, S. F., Shawver, J., Espinosa Morales, A., Salahuddin, H., Tietjen, G., Lindstrom, D., ... Jumaa, M. A. (2017). Stroke care: Initial data from a county-based bypass protocol for patients with acute stroke. *Journal of Neurointerventional Surgery*, *9*(7), 631–635. <https://doi.org/10.1136/neurintsurg-2016-012476>.
- Mazyra, M. V., Berglund, A., Ahmed, N., von Euler, M., Holmin, S., Laska, A. C., ... Eriksson, E. E. (2020). Implementation of a prehospital stroke triage system using symptom severity and teleconsultation in the Stockholm Stroke Triage Study. *JAMA Neurology*, *77*(6), 691–699. <https://doi.org/10.1001/jamaneurol.2020.0319>.
- Duvekot, M. H. C., Venema, E., Rozeman, A. D., Moudrouts, W., Vermeij, F. H., Biekart, M., ... Presto investigators (2021). Comparison of eight prehospital stroke scales to detect intracranial large-vessel occlusion in suspected stroke (PRESTO): A prospective observational study. *The Lancet Neurology*, *20*(3), 213–221. [https://doi.org/10.1016/S1474-4422\(20\)30439-7](https://doi.org/10.1016/S1474-4422(20)30439-7).
- Nguyen, T. T. M., van den Wijngaard, I. R., Bosch, J., van Belle, E., van Zwet, E. W., Dofferhoff-Vermeulen, T., ... Kruyt, N. D. (2021). Comparison of prehospital scales for predicting large anterior vessel occlusion in the ambulance setting. *JAMA Neurology*, *78*(2), 157–164. <https://doi.org/10.1001/jamaneurol.2020.4418>.
- Nazliel, B., Starkman, S., Liebeskind, D. S., Ovbiagele, B., Kim, D., Sanossian, N., ... Saver, J. L. (2008). A brief prehospital stroke severity scale identifies ischemic stroke patients harboring persisting large arterial occlusions. *Stroke*, *39*(8), 2264–2267. <https://doi.org/10.1161/STROKEAHA.107.508127>.
- Schlechtriemen, T., Becker, M., & Hoor, L. (2019). Prehospital management of acute stroke: Using the example of Saarland ambulance services. *Radiologe*, *59*(7), 596–602. <https://doi.org/10.1007/s00117-019-0550-5>.
- Mullen, M. T., Pajeroski, W., Messe, S. R., Mechem, C. C., Jia, J., Abboud, M., ... Band, R. (2018). Geographic modeling to quantify the impact of primary and comprehensive stroke center destination policies. *Stroke*, *49*(4), 1021–1023. <https://doi.org/10.1161/STROKEAHA.118.020691>.
- von Elm, E., Altman, D. G., Egger, M., Pocock, S. J., Gotsche, P. C., Vandenbroucke, J. P., & Initiative, S. (2007). The strengthening of reporting of observational studies in epidemiology (STROBE) statement: Guidelines for reporting observational studies. *The Lancet*, *370*(9596), 1453–1457. [https://doi.org/10.1016/S0140-6736\(07\)61602-X](https://doi.org/10.1016/S0140-6736(07)61602-X).
- Eyding, J., Bartig, D., Weber, R., Katsanos, A. H., Weimar, C., Hacke, W., & Krogias, C. (2019). Inpatient TIA and stroke care in adult patients in Germany - retrospective analysis of nationwide administrative data sets of 2011 to 2017. *Neurological Research and Practice*, *1*(1), 39. <https://doi.org/10.1186/s42466-019-0044-y>.
- Hemphill 3rd, J. C., Greenberg, S. M., Anderson, C. S., Becker, K., Bendok, B. R., Cushman, M., ... Council on Clinical Cardiology (2015). Guidelines for the Management of Spontaneous Intracerebral Hemorrhage: A guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*, *46*(7), 2032–2060. <https://doi.org/10.1161/STR.0000000000000069>.
- Smith, E. E., Kent, D. M., Bulsara, K. R., Leung, L. Y., Lichtman, J. H., Reeves, M. J., ... American Heart Association Stroke Council (2018). Accuracy of prediction instruments for diagnosing large vessel occlusion in individuals with suspected stroke: A systematic review for the 2018 guidelines for the early management of patients with acute ischemic stroke. *Stroke*, *49*(3), e111–e122. <https://doi.org/10.1161/STR.0000000000000160>.
- Turc, G., Maier, B., Naggara, O., Seners, P., Isabel, C., Tisserand, M., ... Oppenheim, C. (2016). Clinical scales do not reliably identify acute ischemic stroke patients with large-artery occlusion. *Stroke*, *47*(6), 1466–1472. <https://doi.org/10.1161/STROKEAHA.116.013144>.
- Zhao, H., Pesavento, L., Coote, S., Rodrigues, E., Salvaris, P., Smith, K., ... Campbell, B. C. V. (2018). Ambulance clinical triage for acute stroke treatment: Paramedic triage algorithm for large vessel occlusion. *Stroke*, *49*(4), 945–951. <https://doi.org/10.1161/STROKEAHA.117.019307>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.